

Reading is the second part of the IELTS test, and takes 60 minutes. It consists of three or sometimes four reading passages of increasing difficulty, and there is a total of 40 questions to answer. Though you can mark and write on the Question Paper, you must enter your answers on the Reading Answer Sheet, and be aware that no extra time is given for transferring your answers from the test booklet to the Reading Answer Sheet.

The Academic and General Training Reading Tests are graded to the same level. However, because the texts in the Academic Reading Test are more challenging overall than those in the General Training Test, more questions need to be answered correctly on a General Training Test to receive the same grade.

IELTS Academic Reading Sample 5 - The Risks of Cigarette Smoke

Last Updated: Sunday, 31 July 2016 13:52

Written by IELTS Mentor

Hits: 34916

You should spend about 20 minutes on Questions 15-28 which are based on Reading Passage 5 below.

The Risks of Cigarette Smoke

Discovered in the early 1800s and named nicotianine, the oily essence now called nicotine is the main active ingredient of tobacco. Nicotine, however, is only a small component of cigarette smoke, which contains more than 4,700 chemical compounds, including 43 cancer-causing substances. In recent times, scientific research has been providing evidence that, years of cigarette smoking vastly increases the risk of developing fatal medical conditions.

In addition to being responsible for more than 85 per cent of lung cancers, smoking is associated with cancers of, amongst others, the mouth, stomach and kidneys, and is thought to cause about 14 percent of leukaemia and cervical cancers. In 1990, smoking caused more than 84,000 deaths, mainly resulting from such problems as pneumonia, bronchitis and influenza. Smoking, it is believed, is responsible for 30 percent of all deaths from cancer and clearly represents the most important preventable cause of cancer in countries like the United States today.

Passive smoking, the breathing in of the side-stream smoke from the burning of tobacco between puffs or of the smoke exhaled by a smoker, also causes a serious health risk. A report published in 1992 by the US Environmental Protection Agency (EPA) emphasised the health dangers, especially from side-stream smoke. This type of smoke contains more, smaller particles and is therefore, more likely to be deposited deep in the lungs. On the basis of this report, the EPA has classified environmental tobacco smoke in the highest risk category for causing cancer.

As an illustration of the health risks, in the case of a married couple where one partner is a smoker and one a non-smoker, the latter is believed to have a 30 percent higher risk of death from heart disease because of passive smoking. The risk of lung cancer also increases over the years of exposure and the figure jumps to 80 percent if the spouse has been smoking four packs a day for 20 years. It has been calculated that 17 percent of cases of lung cancer can be attributed to high levels of exposure to second-hand tobacco smoke during childhood and adolescence.

A more recent study by researchers at the University of California at San Francisco (UCSF) has shown that second-hand cigarette smoke does more harm to non-smokers than to smokers. Leaving aside the philosophical question of whether anyone should have to breathe someone else's cigarette smoke, the report suggests that the smoke experienced by many people in their daily lives is enough to produce substantial adverse effects on a person's heart and lungs.

The report, published in the Journal of the American Medical Association (AMA), was based on the researchers' own earlier research but also includes a review of studies over the past few years. The American Medical Association represents about half of all US doctors and is a strong opponent of smoking. The study suggests that people who smoke cigarettes are continually damaging their cardiovascular system, which adapts in order to compensate for the effects of smoking. It further states that people who do not smoke do not have the benefit of their system adapting to the smoke inhalation. Consequently, the effects of passive smoking are far greater on non-smokers than on smokers.

This report emphasises that cancer is not caused by a single element in cigarette smoke; harmful effects to health are caused by many components. Carbon monoxide, for example, competes with oxygen in red blood cells and interferes with the blood's ability to deliver life-giving oxygen to the heart. Nicotine and other toxins in cigarette smoke activate small blood cells called platelets, which increases the likelihood of blood clots, thereby affecting blood circulation throughout the body.

The researchers criticise the practice of some scientific consultants who work with the tobacco industry for assuming that cigarette smoke has the same impact on smokers as it does on non-smokers. They argue that those scientists are underestimating the damage done by passive smoking and, in support of their recent findings, cite some previous research which points to passive smoking as the cause for between 30,000 and 60,000 deaths from heart attacks each year in the United States. This means that passive smoking is the third most preventable cause of death after active smoking and alcohol-related diseases.

The study argues that the type of action needed against passive smoking should be similar to that being taken against illegal drugs and AIDS (SIDA). The UCSF researchers maintain that the simplest and most cost-effective action is to establish smoke-free workplaces, schools and public places.

Questions 15-17

Choose the appropriate letters **A - D** and write them in boxes **15 -17** on your answer sheet.

15 According to information in the text, leukaemia and pneumonia

- A are responsible for 84,000 deaths each year.
- B are strongly linked to cigarette smoking.
- C are strongly linked to lung cancer.
- D result in 30 percent of deaths per year.

16 According to information in the text, intake of carbon monoxide

- A inhibits the flow of oxygen to the heart.
- B increases absorption of other smoke particles.
- C inhibits red blood cell formation.
- D promotes nicotine absorption.

17 According to information in the text, intake of nicotine encourages

- A blood circulation through the body.
- B activity of other toxins in the blood.
- C formation of blood clots.
- D an increase of platelets in the blood.

Questions 18-21

Do the following statements reflect the claims of the writer in Reading Passage 5? In boxes 18-21 on your answer sheet write:

- YES** if the statement reflects the claims of the writer
- NO** if the statement contradicts the claims of the writer
- NOT GIVEN** if it is impossible to say what the writer thinks about this

- 18 Thirty percent of deaths in the United States is caused by smoking-related diseases.
- 19 If one partner in a marriage smokes, the other is likely to take up smoking.
- 20 Teenagers whose parents smoke are at risk of getting lung cancer at some time during their lives.
- 21 Opponents of smoking financed the UCSF study.

Questions 22-24

Choose ONE phrase from the list of phrases **A - J** below to complete each of the following sentences (Questions **22-24**).

Write the appropriate letters in boxes **22 - 24** on your answer sheet.

- 22 Passive smoking
- 23 Compared with a non-smoker, a smoker
- 24 The American Medical Association

- A** includes reviews of studies in its reports.
- B** argues for stronger action against smoking in public places.
- C** is one of the two most preventable causes of death.
- D** is more likely to be at risk from passive smoking diseases.
- E** is more harmful to non-smokers than to smokers.
- F** is less likely to be at risk of contracting lung cancer.
- G** is more likely to be at risk of contracting various cancers.
- H** opposes smoking and publishes research on the subject.
- I** is just as harmful to smokers as it is to non-smokers.
- J** reduces the quantity of blood flowing around the body.

Questions 25-28

Classify the following statements as being

- A** a finding of the UCSF study
- B** an opinion of the UCSF study

- C** a finding of the EPA report
- D** an assumption of consultants to the tobacco industry

Write the appropriate letters **A—D** in boxes **25—28** on your answer sheet.

NB You may use any letter more than once.

- 25 Smokers' cardiovascular systems adapt to the intake of environmental smoke.
- 26 There is a philosophical question as to whether people should have to inhale others' smoke.
- 27 Smoke-free public places offer the best solution.
- 28 The intake of side-stream smoke is more harmful than smoke exhaled by a smoker.

Click the button to Show/ Hide Answers

Answer:

15. B 16. A 17. C 18. NO 19. NOT GIVEN 20. YES 21. NOT GIVEN 22. E 23. G 24. H 25. A 26. B 27. B 28. C

For more, please visit <http://EngineerMohsinAli.blogspot.com> [This blog is about the hub of Engineering and education or visit here <http://IELTSdocument.weebly.com> for IELTS related materials free of cost.